

Universal Yang–Mills Action on Four Dimensional Manifolds

Kazuyuki FUJII ^{*}, Hiroshi OIKE [†] and Tatsuo SUZUKI [‡]

^{*}Department of Mathematical Sciences

Yokohama City University

Yokohama, 236–0027

Japan

[†]Takado 85–5, Yamagata, 990–2464

Japan

[‡]Department of Mathematical Sciences

Waseda University

Tokyo, 169–8555

Japan

Abstract

The usual action of Yang–Mills theory is given by the quadratic form of curvatures of a principal G bundle defined on four dimensional manifolds. The non–linear generalization which is known as the Born–Infeld action has been given.

^{*}E-mail address : fujii@yokohama-cu.ac.jp

[†]E-mail address : oike@tea.ocn.ne.jp

[‡]E-mail address : suzukita@gm.math.waseda.ac.jp

In this paper we give another non-linear generalization on four dimensional manifolds and call it a universal Yang-Mills action.

The advantage of our model is that the action splits **automatically** into two parts consisting of self-dual and anti-self-dual directions. Namely, we have automatically the self-dual and anti-self-dual equations without solving the equations of motion as in a usual case.

Our method may be applicable to recent non-commutative Yang-Mills theories studied widely.

1 Introduction

The purpose of this paper is to develop a universal Yang-Mills theory on four dimensional manifolds.

The Yang-Mills theory [1] is a fundamental one in elementary particle physics and there is nothing added furthermore. See also [2] and [3] as a part of history. We are interested in some non-linear generalizations of it. One of them is well-known as the Born-Infeld theory [4].

However, we have a dissatisfaction at the model and so we present another non-linear generalization. Our action is related to a Chern character for a curvature matrix defined in the text, while the Born-Infeld one is related to a Chern class for it. Concerning a brief introduction of the Chern theory [5] is recommended.

In the Yang-Mills theory instantons and anti-instantons [6] which are solutions of the self-dual and anti-self-dual equations play a central role.

The typical characteristic of our model is that the action splits **automatically** into two parts consisting of self-dual and anti-self-dual directions without solving the equations of motion as in a usual case. We hope that our theory may change a framework of unified gauge theory.

Although our method in this paper is restricted to four dimensional manifolds because of some technical reasons, it must be extended to higher dimensional cases (maybe, $4n$ -dimensional manifolds). Moreover, it may be applicable to non-commutative Yang-Mills the-

ories studied vigorously, see for example [7], [8]. However, it is beyond our scope in the paper. Further study will be required.

2 Mathematical Preliminaries

We make brief mathematical preliminaries within our necessity for the latter convenience. See [5] for more detailed descriptions. In the following we consider the $\mathbf{C}^\infty$ category, namely $\mathbf{C}^\infty$ -manifolds, $\mathbf{C}^\infty$ -maps, etc.

Let M be a four dimensional manifold and G a classical group, in particular, $U(1)$ and $SU(2)$. We denote by $\{G, P, \pi, M\}$ a principal G bundle on M

$$\pi : P \longrightarrow M, \quad \pi^{-1}(m) \cong G.$$

We consider a theory of connections of the principal G bundle, so let $\mathfrak{g}$ be the Lie algebra of the group G . In the following we treat it locally, which is enough for our purpose. Let U be any open set in M , then a connection $\{A_\mu\}$ (a gauge potential) is

$$A_\mu : U \longrightarrow \mathfrak{g}$$

and the corresponding curvature $\{F_{\mu\nu}\}$ is given by

$$F_{\mu\nu} = \frac{\partial A_\nu}{\partial x_\mu} - \frac{\partial A_\mu}{\partial x_\nu} + [A_\mu, A_\nu] \equiv \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

We note that

$$F_{\mu\nu} : U \longrightarrow \mathfrak{g}, \quad F_{\nu\mu} = -F_{\mu\nu}.$$

For a map $\phi : U \longrightarrow G$, a gauge transformation by ϕ is defined by

$$A_\mu \longrightarrow \phi^{-1} A_\mu \phi + \phi^{-1} \partial_\mu \phi$$

and the curvature is then transformed like

$$F_{\mu\nu} \longrightarrow \phi^{-1} F_{\mu\nu} \phi.$$

3 Universal Yang–Mills Action : Abelian Case

In this section we consider the abelian case $G = U(1)$, which gives us a prototype. For that let us define a curvature matrix

$$\mathcal{F} = \begin{pmatrix} 0 & F_{12} & F_{13} & F_{14} \\ -F_{12} & 0 & F_{23} & F_{24} \\ -F_{13} & -F_{23} & 0 & F_{34} \\ -F_{14} & -F_{24} & -F_{34} & 0 \end{pmatrix}. \quad (1)$$

The abelian Yang–Mills action $\mathcal{A}_{YM} ([1])$ ¹ is given by

$$\mathcal{A}_{YM} \equiv \frac{1}{2} \text{tr} (g\mathcal{F})^2 = -g^2 \sum_{i < j} F_{ij}^2 \quad (2)$$

where g is a coupling constant.

This model is very well-known and there is nothing added furthermore. We are interested in some non-linear generalization(s) of it.

A non-linear extension of the abelian Yang–Mills is known as the Born–Infeld theory ([4]) whose action $\mathcal{A}_{BI}$ is given by

$$\mathcal{A}_{BI} \equiv \sqrt{\det (\mathbf{1}_4 + g\mathcal{F})} \quad (3)$$

where g is a coupling constant.

As to this model and its generalizations to non-abelian groups there are a lot of papers. However, we don't make a comment in this paper.

Since we have a dissatisfaction at the model we present another non-linear extension. Its action $\mathcal{A}$ is

$$\mathcal{A}_{FOS} \equiv \text{tr} e^{g\mathcal{F}} \quad (4)$$

where g is a coupling constant². We want to call it a universal Yang–Mills theory

¹it may be suitable to call it the Maxwell action

²we can completely determine $e^{g\mathcal{F}}$ itself, see [9]

Let us calculate the right hand side of (4). For that we must calculate the characteristic polynomial of $g\mathcal{F}$. It is not difficult to show

$$\begin{aligned}
0 &= |\lambda E - g\mathcal{F}| \\
&= \begin{vmatrix} \lambda & -gF_{12} & -gF_{13} & -gF_{14} \\ gF_{12} & \lambda & -gF_{23} & -gF_{24} \\ gF_{13} & gF_{23} & \lambda & -gF_{34} \\ gF_{14} & gF_{24} & gF_{34} & \lambda \end{vmatrix} \\
&= \lambda^4 + g^2 (F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2) \lambda^2 + g^4 (F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23})^2. \quad (5)
\end{aligned}$$

We set $t = \lambda^2$ for simplicity. Then

$$t^2 + g^2 (F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2) t + g^4 (F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23})^2 = 0.$$

Next let us calculate the discriminant.

$$\begin{aligned}
D &= g^4 (F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2)^2 - 4g^4 (F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23})^2 \\
&= g^4 \left\{ F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2 - 2(F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23}) \right\} \times \\
&\quad \left\{ F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2 + 2(F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23}) \right\} \\
&= g^4 \left\{ (F_{12} - F_{34})^2 + (F_{13} + F_{24})^2 + (F_{14} - F_{23})^2 \right\} \times \\
&\quad \left\{ (F_{12} + F_{34})^2 + (F_{13} - F_{24})^2 + (F_{14} + F_{23})^2 \right\}.
\end{aligned}$$

If we set

$$X_{sd}^2 = (F_{12} - F_{34})^2 + (F_{13} + F_{24})^2 + (F_{14} - F_{23})^2, \quad (6)$$

$$X_{asd}^2 = (F_{12} + F_{34})^2 + (F_{13} - F_{24})^2 + (F_{14} + F_{23})^2 \quad (7)$$

then

$$D = g^4 X_{sd}^2 X_{asd}^2.$$

It is easy to see

$$X_{asd}^2 + X_{sd}^2 = 2 (F_{12}^2 + F_{13}^2 + F_{14}^2 + F_{23}^2 + F_{24}^2 + F_{34}^2),$$

$$X_{asd}^2 - X_{sd}^2 = 4 (F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23}),$$

so

$$t^2 + g^2 \frac{X_{sd}^2 + X_{asd}^2}{2} t + g^4 \frac{(X_{sd}^2 - X_{asd}^2)^2}{16} = 0.$$

From this we have

$$\begin{aligned} t_{\pm} &= g^2 \left\{ -\frac{X_{sd}^2 + X_{asd}^2}{4} \pm \frac{X_{sd}X_{asd}}{2} \right\} = \frac{g^2}{4} \left\{ -(X_{sd}^2 + X_{asd}^2) \pm 2X_{sd}X_{asd} \right\} \\ &= -\frac{g^2}{4} \left\{ (X_{sd}^2 + X_{asd}^2) \mp 2X_{sd}X_{asd} \right\} = -\frac{g^2}{4} (X_{sd} \mp X_{asd})^2, \end{aligned}$$

therefore

$$\begin{aligned} \lambda_1 &= i\frac{g}{2}(X_{sd} + X_{asd}), & \lambda_2 &= -i\frac{g}{2}(X_{sd} + X_{asd}), \\ \lambda_3 &= i\frac{g}{2}(X_{sd} - X_{asd}), & \lambda_4 &= -i\frac{g}{2}(X_{sd} - X_{asd}) \end{aligned} \quad (8)$$

because of $\lambda^2 = t$.

As a result we obtain

$$\begin{aligned} \mathcal{A}_{FOS} &\equiv \text{tr } e^{g\mathcal{F}} \\ &= e^{\lambda_1} + e^{\lambda_2} + e^{\lambda_3} + e^{\lambda_4} \\ &= 2 \cos \left(g \frac{X_{sd} + X_{asd}}{2} \right) + 2 \cos \left(g \frac{X_{sd} - X_{asd}}{2} \right) \\ &= 4 \cos \left(g \frac{X_{sd}}{2} \right) \cos \left(g \frac{X_{asd}}{2} \right). \end{aligned} \quad (9)$$

It is very notable that

$$X_{sd} = 0 \iff F_{12} = F_{34}, \quad F_{13} = -F_{24}, \quad F_{14} = F_{23} \iff \{F_{ij}\} \text{ is } \mathbf{self-dual}, \quad (10)$$

$$X_{asd} = 0 \iff F_{12} = -F_{34}, \quad F_{13} = F_{24}, \quad F_{14} = -F_{23} \iff \{F_{ij}\} \text{ is } \mathbf{anti-self-dual}. \quad (11)$$

The characteristic of our model is that the action (9) splits **automatically** into two parts consisting of self-dual and anti-self-dual directions. Namely, we have **automatically** the self-dual and anti-self-dual equations without solving the equations of motion as in a usual case.

Last in this section let us make an important comment on the action (9). Since the target (as a map) of curvatures $\{F_{ij}\}$ is the Lie algebra $u(1) = \sqrt{-1}\mathbf{R}$ it may be appropriate to write $F_{ij} = \sqrt{-1}G_{ij}$. Then (9) is changed into the more suitable form

$$\mathcal{A}_{FOS} = 4 \cosh\left(g\frac{Y_{sd}}{2}\right) \cosh\left(g\frac{Y_{asd}}{2}\right) \quad (12)$$

with

$$Y_{sd}^2 = (G_{12} - G_{34})^2 + (G_{13} + G_{24})^2 + (G_{14} - G_{23})^2 \geq 0, \quad (13)$$

$$Y_{asd}^2 = (G_{12} + G_{34})^2 + (G_{13} - G_{24})^2 + (G_{14} + G_{23})^2 \geq 0. \quad (14)$$

The same thing will hold for **non-abelian cases** in the following.

4 Universal Yang–Mills Action : Non–Abelian Case

In this section we consider the non-abelian case $G = SU(2)$. From the form in (9) it is natural to conjecture

$$\mathcal{A}_{FOS} = 4 \operatorname{tr} \cos\left(g\frac{X_{sd}}{2}\right) \cos\left(g\frac{X_{asd}}{2}\right) \quad (15)$$

where

$$X_{sd}^2 = (F_{12} - F_{34})^2 + (F_{13} + F_{24})^2 + (F_{14} - F_{23})^2, \quad (16)$$

$$X_{asd}^2 = (F_{12} + F_{34})^2 + (F_{13} - F_{24})^2 + (F_{14} + F_{23})^2. \quad (17)$$

However, a natural question arises. What is a curvature matrix giving (15) as a result ? This is not an easy problem.

As a first trial we generalize (1) as

$$\mathcal{F} = \begin{pmatrix} \mathbf{0}_2 & F_{12} & F_{13} & F_{14} \\ -F_{12} & \mathbf{0}_2 & F_{23} & F_{24} \\ -F_{13} & -F_{23} & \mathbf{0}_2 & F_{34} \\ -F_{14} & -F_{24} & -F_{34} & \mathbf{0}_2 \end{pmatrix} \quad (18)$$

where

$$F_{ij} = F_{ij}^1 \sigma_1 + F_{ij}^2 \sigma_2 + F_{ij}^3 \sigma_3 : U \subset M \longrightarrow su(2) \quad (19)$$

with the Pauli matrices $\{\sigma_1, \sigma_2, \sigma_3\}$ defined by

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (20)$$

In the following it is better to use

$$F_{ij} \longrightarrow \hat{F}_{ij} = \begin{pmatrix} F_{ij}^1 \\ F_{ij}^2 \\ F_{ij}^3 \end{pmatrix}, \quad (21)$$

so we set for simplicity

$$\mathbf{a} \equiv \hat{F}_{12}, \quad \mathbf{b} \equiv \hat{F}_{13}, \quad \mathbf{c} \equiv \hat{F}_{14}, \quad \mathbf{d} \equiv \hat{F}_{23}, \quad \mathbf{e} \equiv \hat{F}_{24}, \quad \mathbf{f} \equiv \hat{F}_{34}. \quad (22)$$

Here we prepare the notations : for three three-dimensional vectors $\mathbf{x}, \mathbf{y}, \mathbf{z}$ we set the inner products, exterior products and scalar triple product respectively as

$$(\mathbf{xy}) \equiv \mathbf{x} \cdot \mathbf{y}, \quad (\mathbf{xx}) \equiv \mathbf{x}^2, \quad (\mathbf{yy}) \equiv \mathbf{y}^2,$$

$$[\mathbf{xy}] \equiv \mathbf{x} \wedge \mathbf{y}, \quad \text{etc},$$

$$|\mathbf{xyz}| \equiv (\mathbf{x} [\mathbf{yz}]) = \mathbf{x} \cdot \mathbf{y} \wedge \mathbf{z} = |\mathbf{x}, \mathbf{y}, \mathbf{z}|.$$

Now we write down a useful relation : for

$$A = a^1 \sigma_1 + a^2 \sigma_2 + a^3 \sigma_3 \implies \mathbf{a} = (a^1, a^2, a^3)^T$$

we have

$$\begin{aligned} A^2 &= \left\{ (a^1)^2 + (a^2)^2 + (a^3)^2 \right\} \mathbf{1}_2 + a^1 a^2 \{\sigma_1, \sigma_2\} + a^1 a^3 \{\sigma_1, \sigma_3\} + a^2 a^3 \{\sigma_2, \sigma_3\} \\ &= \left\{ (a^1)^2 + (a^2)^2 + (a^3)^2 \right\} \mathbf{1}_2 = \mathbf{a}^2 \mathbf{1}_2. \end{aligned} \quad (23)$$

By way of trial we define the universal Yang–Mills action $\mathcal{A}$ like

$$\mathcal{A}_{FOS} \equiv \text{Tr } e^{g\mathcal{F}} \quad (24)$$

with (18), where g is a coupling constant. Here, to avoid a confusion we used the notation Tr not tr in (4) because each component of $\mathcal{F}$ in (18) is a matrix.

Let us calculate the right hand side of (24), which is hard task this time. The characteristic polynomial of $g\mathcal{F}$ is

$$\begin{aligned}
0 &= |\lambda E - g\mathcal{F}| \\
&= \begin{vmatrix} \lambda \mathbf{1}_2 & -gF_{12} & -gF_{13} & -gF_{14} \\ gF_{12} & \lambda \mathbf{1}_2 & -gF_{23} & -gF_{24} \\ gF_{13} & gF_{23} & \lambda \mathbf{1}_2 & -gF_{34} \\ gF_{14} & gF_{24} & gF_{34} & \lambda \mathbf{1}_2 \end{vmatrix} \\
&= \left[\lambda^4 + g^2 (\mathbf{a}^2 + \mathbf{b}^2 + \mathbf{c}^2 + \mathbf{d}^2 + \mathbf{e}^2 + \mathbf{f}^2) \lambda^2 - 2ig^3 (|\mathbf{abd}| + |\mathbf{ace}| + |\mathbf{bcf}| + |\mathbf{def}|) \lambda + \right. \\
&\quad \left. g^4 \text{PF}(\mathcal{F}) \right]^2
\end{aligned} \tag{25}$$

where $\text{PF}(\mathcal{F})$ is the Pfaffian of $\mathcal{F}$ and is given by

$$\begin{aligned}
\text{PF}(\mathcal{F}) &= \mathbf{a}^2 \mathbf{f}^2 + \mathbf{b}^2 \mathbf{e}^2 + \mathbf{c}^2 \mathbf{d}^2 - 2 \{ (\mathbf{ab})(\mathbf{ef}) - (\mathbf{af})(\mathbf{be}) + (\mathbf{ae})(\mathbf{bf}) - (\mathbf{ac})(\mathbf{df}) + \\
&\quad (\mathbf{af})(\mathbf{cd}) - (\mathbf{ad})(\mathbf{cf}) + (\mathbf{bc})(\mathbf{de}) - (\mathbf{be})(\mathbf{cd}) + (\mathbf{bd})(\mathbf{ce}) \}.
\end{aligned} \tag{26}$$

In the calculation the relation (23) plays an important role.

Here by neglecting the λ -term³ we have

$$t^2 + g^2 (\mathbf{a}^2 + \mathbf{b}^2 + \mathbf{c}^2 + \mathbf{d}^2 + \mathbf{e}^2 + \mathbf{f}^2) t + g^4 \text{PF}(\mathcal{F}) = 0$$

for $t = \lambda^2$. Let us calculate the discriminant

$$\begin{aligned}
D &= g^4 (\mathbf{a}^2 + \mathbf{b}^2 + \mathbf{c}^2 + \mathbf{d}^2 + \mathbf{e}^2 + \mathbf{f}^2)^2 - 4g^4 \text{PF}(\mathcal{F})^2 \\
&= g^4 \left\{ \mathbf{a}^2 + \mathbf{b}^2 + \mathbf{c}^2 + \mathbf{d}^2 + \mathbf{e}^2 + \mathbf{f}^2 + 2\text{PF}(\mathcal{F}) \right\} \times \\
&\quad \left\{ \mathbf{a}^2 + \mathbf{b}^2 + \mathbf{c}^2 + \mathbf{d}^2 + \mathbf{e}^2 + \mathbf{f}^2 - 2\text{PF}(\mathcal{F}) \right\}.
\end{aligned}$$

³we can formally solve (25) by use of the formula given by Ferrari or Euler, see for example [10]

Here we define

$$\tilde{X}_{sd}^2 = \{(\mathbf{a} - \mathbf{f})^2 + (\mathbf{b} + \mathbf{e})^2 + (\mathbf{c} - \mathbf{d})^2\}, \quad (27)$$

$$\tilde{X}_{asd}^2 = \{(\mathbf{a} + \mathbf{f})^2 + (\mathbf{b} - \mathbf{e})^2 + (\mathbf{c} + \mathbf{d})^2\} \quad (28)$$

for the latter convenience. For

$$\tilde{X}_{sd}^2 \tilde{X}_{asd}^2 = \{(\mathbf{a} - \mathbf{f})^2 + (\mathbf{b} + \mathbf{e})^2 + (\mathbf{c} - \mathbf{d})^2\} \{(\mathbf{a} + \mathbf{f})^2 + (\mathbf{b} - \mathbf{e})^2 + (\mathbf{c} + \mathbf{d})^2\}$$

a long calculation leads to

$$\begin{aligned} D - g^4 \tilde{X}_{sd}^2 \tilde{X}_{asd}^2 = & -4g^4 \left[([\mathbf{af}] - [\mathbf{be}] + [\mathbf{cd}])^2 \right. \\ & \left. - 4 \{([\mathbf{ab}][\mathbf{ef}] - ([\mathbf{ac}][\mathbf{df}] + ([\mathbf{bc}][\mathbf{de}]))\} \right]. \end{aligned} \quad (29)$$

Therefore we have

$$D = g^4 \tilde{X}_{sd}^2 \tilde{X}_{asd}^2 \mod \{\text{Exterior Products}\}. \quad (30)$$

Since

$$|\mathbf{abd}| + |\mathbf{ace}| + |\mathbf{bcf}| + |\mathbf{def}| = 0 \mod \{\text{Exterior Products}\},$$

the equation

$$0 = |\lambda E - g\mathcal{F}| \mod \{\text{Exterior Products}\} \quad (31)$$

gives

$$t^2 + g^2 \frac{\tilde{X}_{sd}^2 + \tilde{X}_{asd}^2}{2} t + g^4 \frac{(\tilde{X}_{asd}^2 - \tilde{X}_{sd}^2)^2}{16} = 0$$

and so

$$t_{\pm} = -\frac{g^2}{4} (\tilde{X}_{sd} \mp \tilde{X}_{asd})^2.$$

Therefore

$$\begin{aligned} \lambda_1 &= i\frac{g}{2}(\tilde{X}_{sd} + \tilde{X}_{asd}), & \lambda_2 &= -i\frac{g}{2}(\tilde{X}_{sd} + \tilde{X}_{asd}), \\ \lambda_3 &= i\frac{g}{2}(\tilde{X}_{sd} - \tilde{X}_{asd}), & \lambda_4 &= -i\frac{g}{2}(\tilde{X}_{sd} - \tilde{X}_{asd}). \end{aligned} \quad (32)$$

That is, the solutions are $\{\lambda_1, \lambda_1, \lambda_2, \lambda_2, \lambda_3, \lambda_3, \lambda_4, \lambda_4\}$ because they are two-fold.

Finally we obtain the desired result

$$\begin{aligned}
\mathcal{A}_{FOS} &\equiv \text{Tr } e^{g\mathcal{F}} \mod \{\text{Exterior Products}\} \\
&= 2 \left(e^{\lambda_1} + e^{\lambda_2} + e^{\lambda_4} + e^{\lambda_4} \right) \\
&= 8 \cos \left(g \frac{\tilde{X}_{sd}}{2} \right) \cos \left(g \frac{\tilde{X}_{asd}}{2} \right) \\
&= 4 \text{tr} \cos \left(g \frac{X_{sd}}{2} \right) \cos \left(g \frac{X_{asd}}{2} \right). \tag{33}
\end{aligned}$$

A comment is in order. We would like to point out a similarity between our idea and a topic in Topology. The first homology group $H_1(M, \mathbf{Z})$ of M is an abelian group, while the first homotopy group (fundamental group) $\pi_1(M)$ of M is in general non-abelian. However, there is a famous relation

$$H_1(M, \mathbf{Z}) \cong \pi_1(M) / [\pi_1(M), \pi_1(M)] = \pi_1(M) \mod [\pi_1(M), \pi_1(M)]$$

where $[\pi_1(M), \pi_1(M)]$ is the commutator subgroup generated by $\{aba^{-1}b^{-1} \mid a, b \in \pi_1(M)\}$. See for example [5]. Our method is **a kind of abelianization**.

It should be noted that the gauge groups of our model are restricted to $U(1)$ and $SU(2)$, which is of course not enough and we must treat the general $SU(n)$ for $n \in \mathbf{N}$. It is natural to conjecture

$$\mathcal{A}_{FOS} = 4 \text{tr} \cos \left(g \frac{X_{sd}}{2} \right) \cos \left(g \frac{X_{asd}}{2} \right)$$

with

$$\begin{aligned}
F_{ij} &: U \subset M \longrightarrow su(n), \\
X_{sd}^2 &= (F_{12} - F_{34})^2 + (F_{13} + F_{24})^2 + (F_{14} - F_{23})^2, \\
X_{asd}^2 &= (F_{12} + F_{34})^2 + (F_{13} - F_{24})^2 + (F_{14} + F_{23})^2.
\end{aligned}$$

At the present time we don't know how to derive this action from a curvature matrix like (18). We leave it to some readers as a challenging problem.

Our model is also based on four dimensional manifolds M , which is too restrictive in the view point of M–theory or F–theory. We want to generalize M to a $4n$ –dimensional manifold and treat an action like (33).


Moreover, it may be interesting to apply our method to non–commutative Yang–Mills theory (for example [7], [8]), which is however beyond our scope in this paper.

5 Discussion

In this letter we presented the non–linear generalization of the Yang–Mills theory which is different from the Born–Infeld theory. Our action is related to a Chern Character, while that of Born–Infeld is related to a Chern class. Our model has an advantage.

Details of the paper and further developments including some quantization will be published elsewhere [9].

Last let us make a brief comment. We are studying a quantum computation based on Cavity QED whose image is



The general setting for a quantum computation based on Cavity QED :

the dotted line means a single photon inserted in the cavity and

all curves mean external laser fields subjected to atoms

See [11], [12] in detail. Usually it is based on two level system of atoms. However, to take a multi level system of them into consideration may be better from the view point of decoherence which is a severe problem in quantum computation.

It seems that the universal Yang–Mills theory developed in this paper gets on well with a quantum computation based on four level system, see for example [13], [14].

The gauge theory is useful even in quantum computation. This has been pointed out by Fujii [15], [16] in the context of Holonomic Quantum Computation. In a forthcoming paper we want to point out a deeper relation between the gauge theory and some quantum computation (Cavity QED quantum computation if possible).

References

- [1] C. N. Yang and R. L. Mills : Conservation of isotopic spin and isotopic gauge invariance, Phys. Rev. 96 (1954), 191.
- [2] M. Ikeda and Y. Miyachi : On an extended framework for the description of elementary particles, Prog. Theor. Phys. 16 (1956), 537.
- [3] T. T. Wu and C. N. Yang : Concept of nonintegrable phase factors and global formulation of gauge fields, Phys. Rev. D 12 (1975), 3845.
- [4] M. Born and L. Infeld : Foundations of the new field theory, Proc. Royal Soc. (London), A 144 (1934), 425.
- [5] M. Nakahara : Geometry, Topology and Physics, IOP Publishing Ltd, 1990.
- [6] A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Yu. S. Tyupkin : Pseudoparticle solutions of the Yang–Mills equations, Phys. Lett. B 59 (1975), 85.
- [7] A. Connes : Noncommutative Geometry, Academic Press, 1994.
- [8] N. A. Nekrasov : Noncommutative Instantons Revisited, Commun. Math. Phys. 241 (2003) 143, hep-th/0010017.
- [9] K. Fujii, H. Oike and T. Suzuki : in preparation.

- [10] K. Fujii : A Modern Introduction to Cardano and Ferrari Formulas in the Algebraic Equations, quant-ph/0311102.
- [11] K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime, J. Opt. B: Quantum and Semiclass. Opt, 6(2004) 502, quant-ph/0407014.
- [12] K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime II : Complete Construction of the Controlled–Controlled NOT Gate, to appear in the book "Quantum Computing : New Research", 2006, Nova Science Publishers, Inc (USA), quant-ph/0501046.
- [13] K. Fujii : Study on Dynamics of N Level System of Atom by Laser Fields, quant-ph/0512126.
- [14] V. Ramakrishna and H. Zhou : On The Exponential of Matrices in $su(4)$, math-ph/0508018.
- [15] K. Fujii : Note on Coherent States and Adiabatic Connections, Curvatures, J. Math. Phys. 41(2000), 4406, quant-ph/9910069.
- [16] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, J. Applied Math, 2(2002), 371, quant-ph/0103011.